

Electromagnetic Theory: Class Exam II

15 November 2019

Name: _____

Total: /50

Instructions

- There are 4 questions on 7 pages.
- Show your reasoning and calculations and always explain your answers.

Physical constants and useful formulae

Permittivity of free space $\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2$

Permeability of free space $\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2$

Charge of an electron $e = -1.60 \times 10^{-19} \text{ C}$

Integrals

$$\int \sin(ax) \sin(bx) dx = \frac{\sin((a-b)x)}{2(a-b)} - \frac{\sin((a+b)x)}{2(a+b)} \quad \text{if } a \neq b$$

$$\int \cos(ax) \cos(bx) dx = \frac{\sin((a-b)x)}{2(a-b)} + \frac{\sin((a+b)x)}{2(a+b)} \quad \text{if } a \neq b$$

$$\int \sin(ax) \cos(ax) dx = \frac{1}{2a} \sin^2(ax)$$

$$\int \sin^2(ax) dx = \frac{x}{2} - \frac{\sin(2ax)}{4a}$$

$$\int \cos^2(ax) dx = \frac{x}{2} + \frac{\sin(2ax)}{4a}$$

$$\int x \sin^2(ax) dx = \frac{x^2}{4} - \frac{x \sin(2ax)}{4a} - \frac{\cos(2ax)}{8a^2}$$

$$\int x^2 \sin^2(ax) dx = \frac{x^3}{6} - \frac{x^2}{4a} \sin(2ax) - \frac{x}{4a^2} \cos(2ax) + \frac{1}{8a^3} \sin(2ax)$$

Question 1


A spherical conductor of radius R is charged. There is no charge outside of the conductor. Which of the following (choose one) is a possible potential, given in spherical coordinates, at all points **on and beyond** the surface of the conductor?

- i) $V(r, \theta, \phi) = \frac{1}{4\pi\epsilon_0} \frac{Q}{R}$
- ii) $V(r, \theta, \phi) = \frac{1}{4\pi\epsilon_0} \frac{Q}{r}$
- iii) $V(r, \theta, \phi) = \frac{1}{4\pi\epsilon_0} \frac{QR}{r^2}$
- iv) $V(r, \theta, \phi) = \frac{1}{4\pi\epsilon_0} \frac{Q \cos \phi}{r}$
- v) $V(r, \theta, \phi) = \text{constant}$

Briefly explain your answer.

Question 2

An infinitely long cylindrical shell, of negligible thickness and radius b , surrounds an infinitely long solid cylindrical rod, of radius a . Their axes are both along the z -axis and a view down the length of this axis is illustrated. The volume current density in the rod is (in cylindrical coordinates) $\mathbf{J} = \frac{3\alpha}{2\pi a^3} s \hat{\mathbf{z}}$ where α is a constant with the dimensions of current. The current along the cylindrical shell, flows in the $-\hat{\mathbf{z}}$ direction, is uniformly distributed across the surface and is such that the total current flowing down the shell is exactly opposite to that flowing through the cylinder.



- a) Show that the surface current density in the cylindrical shell is

$$\mathbf{K} = -\frac{\alpha}{2\pi b} \hat{\mathbf{z}}.$$

Question 2 continued ...

b) Determine the magnetic field at **all locations**.


/16

Question 3

A circular loop with radius R lies in the xy plane as illustrated and carries a current I counterclockwise. A hidden source current produces the magnetic field

$$\mathbf{B} = \alpha x \hat{\mathbf{z}}$$

where $\alpha > 0$ is a constant. Show qualitatively that the net force exerted by the field on the loop is not zero and determine the net force exerted by the field on the loop.




Question 4

In the following question do **either part a) or part b) for full credit**. If you do both parts, each will be graded and you will be given the highest score that you obtained for one of the parts.

- a) A magnetic dipole has magnetic moment $\mathbf{m} = m\hat{\mathbf{x}}$. Determine the magnetic vector potential in spherical coordinates and use this to determine the magnetic field produced by the dipole in spherical coordinates.

Question 4 continued ...

- b) A loop of radius R carries charge of uniform linear density, λ , and lies in the xy plane. The loop rotates about the axis through its center, as illustrated in the diagram, so that any point moves with speed v . Determine an expression for the magnetic dipole moment of the loop in terms of R, λ, v and constants.



/8