


## Electromagnetic Theory: Homework 14

Due: 15 October 2019

### 1 Conductor with cavities

A conductor contains a spherical cavity with radius  $R$ . The conductor is initially neutral. Subsequently a point particle with charge  $Q$  is placed at the center of the cavity.



- a) Gauss' law can be used to show that for any surface, the charge density on the surface,  $\sigma$ , is given by


$$\mathbf{E}_{\text{above}} - \mathbf{E}_{\text{below}} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{n}}$$

where the electric fields are those immediately above and below the surface and  $\hat{\mathbf{n}}$  is the unit vector perpendicular to the surface pointing in the direction from below to above. Use this to determine the charge density on the inner surface of the cavity.

- b) Determine the total charge on the outer surface of the conductor. Explain your answer.
- c) Can you use your results to determine the charge density on the outer surface? Explain your answer.
- d) Suppose that an additional point charge is placed anywhere outside the conductor. Which of the following stay the same and which change: electrostatic potential inside the cavity, charge density on the cavity, total charge on the outer surface, electric field beyond the conductor? Explain your answer.
- e) Suppose that an additional cavity is created inside the conductor and another charge  $q$  is placed at the center of this. Which of the following stay the same and which change: electrostatic potential inside the original cavity, charge density on the original cavity, total charge on the outer surface? Explain your answer.

### 2 Charge in a conducting shell

A point charge,  $Q$  is placed at the center of a thick conducting shell as illustrated.




- a) Determine the electrostatic potential and electric field inside the shell.
- b) Assuming that the shell was neutral before the point charge was placed at the center, determine the charge densities on both surfaces of the shell.

### 3 Cavity surface charge density

Two conductors each contain a spherical cavity. Each conductor is initially neutral. Subsequently a point particle is placed at the center of the cavity. The radius of the cavity in case A is  $R$  and that of case B is  $2R$ . The particle in the center of the cavity in case A has charge  $Q$  and that of case B has charge  $2Q$ . Let  $\sigma_A$  be the charge density on the inner surface of the cavity in case A and  $\sigma_B$  that of case B. Which of the following is true? Explain your answer.


- i)  $\sigma_A = \frac{1}{4}\sigma_B$ .
- ii)  $\sigma_A = \frac{1}{2}\sigma_B$ .
- iii)  $\sigma_A = \sigma_B$ .
- iv)  $\sigma_A = 2\sigma_B$ .
- v)  $\sigma_A = 4\sigma_B$ .



### 4 Extrema of potentials due to point charges

Two identical point charges are situated as illustrated.

- a) Determine an expression for the electrostatic potential  $V(x, y)$  at all locations.
- b) Use the potential to find locations where the net force on any charged particle is zero.
- c) Describe whether these are local maxima or minima of the potential. Are they stable equilibria?



### 5 Laplace's equation in spherical coordinates

- a) Consider a region in which there is no charge. Write Laplace's equation in terms of spherical coordinates. Solve this assuming that the potential only depends on  $r$ .
- b) What type(s) of charge distribution could give this potential?
- c) Is this potential applicable to the region beyond a charged spherical conductor? Explain your answer.
- d) Is this potential applicable to the region beyond a charged rectangular conductor? Explain your answer.