/70

General Physics: Class Exam I

15 February 2019

Name: _____ Total:

Instructions

• There are 8 questions on 5 pages.

• Show your reasoning and calculations and always explain your answers.

Physical constants and useful formulae

 $e = 1.61 \times 10^{-19} \,\mathrm{C}$ $q_{\text{electron}} = -e \quad q_{\text{proton}} = +e$ $m_{\text{electron}} = 9.11 \times 10^{-31} \,\mathrm{kg}$ $m_{\text{proton}} = 1.67 \times 10^{-27} \,\mathrm{kg}$ $k = 8.99 \times 10^9 \,\mathrm{Nm^2/C^2}$

Question 1

A miniscule sphere contains 2000 excess electrons (the remaining charges cancel out).

- a) Determine the total charge of this sphere.
- b) Determine the magnitude of the force that the sphere exerts on a single electron at a distance of 1.0×10^{-10} m from the sphere.

Two fixed point charged particles are located as illustrated. The charge of particle A is $+4.0 \times 10^{-9}$ C and the charge of particle B is -1.0×10^{-9} C. Determine the net electric field at the location labeled P.



Two identical positively charged particles are held fixed as illustrated. Let E_A be the magnitude of the net electric field at A and E_B be the magnitude of the net electric field at B. Which of the following (choose one) is true?

i) $E_A = E_B$

ii)
$$E_A > E_B$$

iii) $E_A < E_B$

Question 4

Hidden sources produce the illustrated electric field. In separate experiments various charged particles are released from rest, either from point A or else point B.

 a) A negatively charged particle is released from rest at point
B. In which direction will it move immediately after it has been released? Explain your answer.



 $\uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow_{B}$ $\uparrow \uparrow \uparrow \uparrow_{A} \uparrow \uparrow$

/5

b) A particle with charge +8.0 C is placed at point A and the force on it is 16 N. This is replaced by a particle with charge +4.0 C. Determine the force on this new particle.

/10

A distribution of charges produces the illustrated electric potential. An electron moves from the illustrated initial to final points.

- a) Which of the following (choose one) is true for change in electric potential energy of the electron?
 - i) $\Delta U_{\text{elec}} = 0.$
 - ii) $\Delta U_{\text{elec}} > 0.$
 - iii) $\Delta U_{\text{elec}} < 0.$
- b) Which of the following (choose one) is true for the electron?
 - i) Its speed at the final location is the same as at the initial location.
 - ii) Its speed at the final location is larger than at the initial location.
 - iii) Its speed at the final location is smaller than at the initial location.

Question 6

Two charges are held fixed as illustrated. Let $V_A = 2C$ A be the electric potential at point A and V_B be the \bigcirc X potential at B. Which of the following (choose one) is true?

- i) $V_A = V_B$ and both are 0 V.
- ii) $V_A = V_B$ and neither are 0 V.

iii)
$$V_A > V_B$$

iv) $V_A < V_B$.

Question 7

Two parallel plates are held 0.080 m apart. The electric potential difference from one plate to the other is $2.0 \times 10^3 \text{ V}$. Determine the magnitude of the electric field between the plates.



/10

В

X

/5

 $+2\,\mathrm{C}$

 \oplus

An atomic nucleus consists of a number of protons, whose total charge is 8.0×10^{-18} C. Another single proton is released from rest at a distance of 2.0×10^{-10} m from the nucleus.

a) Determine the electric potential produced by the nucleus at the location where the proton is released.

b) Determine the speed of the proton when it is infinitely far from the nucleus.