

Electromagnetic Theory: Class Exam I

2 October 2020

Name: _____

Total: /50

Instructions

- There are 5 questions on 6 pages.
- Show your reasoning and calculations and always explain your answers.

Physical constants and useful formulae

Permittivity of free space $\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2$

Charge of an electron $e = -1.60 \times 10^{-19} \text{ C}$

Integrals

$$\int \sin(ax) \sin(bx) dx = \frac{\sin((a-b)x)}{2(a-b)} - \frac{\sin((a+b)x)}{2(a+b)} \quad \text{if } a \neq b$$

$$\int \cos(ax) \cos(bx) dx = \frac{\sin((a-b)x)}{2(a-b)} + \frac{\sin((a+b)x)}{2(a+b)} \quad \text{if } a \neq b$$

$$\int \sin(ax) \cos(ax) dx = \frac{1}{2a} \sin^2(ax)$$

$$\int \sin^2(ax) dx = \frac{x}{2} - \frac{\sin(2ax)}{4a}$$

$$\int \cos^2(ax) dx = \frac{x}{2} + \frac{\sin(2ax)}{4a}$$

$$\int x \sin^2(ax) dx = \frac{x^2}{4} - \frac{x \sin(2ax)}{4a} - \frac{\cos(2ax)}{8a^2}$$

$$\int x^2 \sin^2(ax) dx = \frac{x^3}{6} - \frac{x^2}{4a} \sin(2ax) - \frac{x}{4a^2} \cos(2ax) + \frac{1}{8a^3} \sin(2ax)$$

Question 1

A solid sphere with radius R has charge density


$$\rho(\mathbf{r}') = \frac{\beta}{r'}$$

where β is a constant with units of C/m^2 . Determine an expression for the electric field at any location inside or outside of the sphere. Express your answer in terms of the total charge on the sphere.

/15

Question 2

Two widely separated spheres carry identical total charges that are uniformly distributed. Sphere A has radius R and sphere B has radius $2R$ where R is some constant. Consider two points (indicated by “o”) that are distance $3R$ from the center of each sphere. How does the magnitude of the electric field for sphere A at this point compare (same, larger, smaller) to that for B at the same point? Explain your answer.



/5

Question 3

The electric field produced by an unknown charge distribution is (in spherical coordinates)

$$\mathbf{E} = E_0 e^{-\lambda r} \hat{\mathbf{r}}$$

where $\lambda > 0$ is a constant with units of m^{-1} and E_0 is a constant with units of N/C .

a) Determine the electrostatic potential at all points, assuming that $V \rightarrow 0$ as $r \rightarrow \infty$.


b) A particle with charge $q > 0$ and mass m is placed at a point r_0 and is released from rest. In which direction does the particle move and what is the maximum speed that it attains?

/12

Question 4

Two infinite slabs are parallel to each other. Each slab has two surfaces and the charge on each surface is uniformly distributed with the illustrated densities ($\sigma > 0$). There is no charge inside the slabs.

- a) Describe as precisely as possible how the magnitude of the electric field inside slab A is related to that of the field inside slab B. Explain your answer.



- b) Suppose that the separation between the slabs was increased. Would the electrostatic potential difference across the gap between the plates increase, decrease or stay constant? Explain your answer.

/8

Question 5

A possible electric field has the form (in cylindrical coordinates)

$$\mathbf{E} = \frac{\alpha}{s} \hat{\phi}$$

where $\alpha > 0$.

a) Sketch the electric field in the xy plane.

b) Determine whether this is a possible electrostatic field.

/10